

RESPONSE OF JAPANESE BUNCHING ONION (*Allium fistulosum* L.) TO NITROGEN FERTILIZATION

Eugeniusz Kołota, Katarzyna Adamczewska-Sowińska,
Cecylia Uklańska-Pusz

Wrocław University of Environmental and Life Sciences

Abstract. New pseudostem cultivars of Japanese bunching onion may be a good alternative to common bulb onion grown for bunches in early spring season. There is still limited knowledge upon the requirement of this vegetable species for nitrogen fertilization. In two factorial field experiment, calcium nitrate, ammonium nitrate and Entec 26 – nitrogen mineral fertilizer containing DMPP nitrification inhibitor used as the source of N and applied at the rates of 75, 150 and 225 kg·ha⁻¹ were compared. Seed propagated transplants produced in the greenhouse were planted in early April and harvested after 10 weeks of cultivation in the open field. The yield and nutritional value of edible parts expressed by the content of dry matter, vitamin C, total chlorophyll, carotenoids, volatile oils, total N, NO₃ – N, P K Ca and Mg were estimated. Results of the study proved that all tested fertilizers were equally valuable sources of nitrogen for Japanese bunching onion. The use of Entec 26 was associated with higher amounts of total chlorophyll and carotenoids in edible part of plants if compared to commonly recommended ammonium nitrate and similar to calcium nitrate. The other important advantage of this fertilizer was a substantial decline of nitrates content. The increment of preplant nitrogen rate from 75 to 150 and 225 kg·ha⁻¹ did not affect the crop yield and significantly enhanced the nitrates accumulation in plants at harvest.

Key words: nitrogen form, N rate, yield, nutritional crop value

INTRODUCTION

Allium species are grown worldwide and are popular in many countries because of high level of phytochemicals that provide health benefits and have specific flavour properties [Griffith et al. 2002]. Among them, onion and leek are the most popular edible *Alliums* grown in most regions of the world [Rabinowitch and Currah 2002], while Japanese bunching onion (*Allium fistulosum* L.), predominantly in Eastern Asia countries.

Corresponding author: Katarzyna Adamczewska-Sowińska, Department of Horticulture, Wrocław University of Environmental and Life Sciences, Pl. Grunwaldzki 24a, 50-363 Wrocław, Poland, tel. (+48) 71 320 17 25, e-mail: katarzyna.a-sowinska@up.wroc.pl

Wild relatives of Japanese bunching onion originated in the region of West China and some neighboring countries of middle Asia, where they have been cultivated for more than 2000 years [Wang et al. 2005, Liu et al. 2009]. It was probably introduced into Europe by the end of Middle Ages receiving the name of Welsh onion [Helm 1956] and mostly grown in home gardens. Although perennial, in Asia the plant is mostly grown as an annual and sometimes as a biennial for its green tops or blanched leeklike pseudostem [Rubatzky and Yamaguchi 1997, Warade and Shinde 1998]. It can be also used in the seedling stage in special Japanese dishes. In this case the plants are harvested when they are about 8–10 cm high [Jones and Mann 1963, Wang et al. 2005].

In Poland, Japanese bunching onion is still a minor crop grown predominantly as a perennial plant in home gardens for the use of green tops in early spring season [Kotlińska and Kojima 2000, Tendaj and Mysiak 2007a, 2010]. New pseudostem type cultivars available nowadays in the market producing little number of tillers [Tendaj and Mysiak 2011] can be grown for the use of whole plants harvested in early growth stages or later, usually in September, with blanched pseudostems like leek species. [Kotlińska and Kojima 2000, Kotlińska et al. 2005]. Such type of cultivars having short growing period can be considered as an alternative green bunching onion to commonly used onion for this purpose [Grevsen 1989].

The advantageous features of Japanese bunching onion are the significant length of white portion of pseudostem, fewer but longer leaves, more erect foliage habit and stronger flavour in comparison to the common onion [Abbey et al. 2002, Lazić et al. 2002, Wang et al. 2005]. Martinez et al. [2005] indicate resistance of *Allium fistulosum* L. to some diseases and pests of bulb onion, and pink root rot among them.

Japanese bunching onion is generally appreciated for its high nutritional value, and unique flavour due to substantial amounts of vitamins, macro- and micronutrients, volatile oils as well as flavonoids possessing high antioxidant properties [He et al. 1989, Horbowicz and Kotlińska 1998, 2000, Kotlińska and Kojima 2000, Lazić et al. 2002, Stainer et al. 2006, Higashio et al. 2007, Tendaj and Mysiak 2007b, Mysiak and Tendaj 2006, 2008].

Growth of bunching onion, its yield and nutritional value of the crop can be highly influenced by the nutrients supply. The recommended rates of fertilizers in cultivation for blanched pseudostems on soil with high organic matter include 200–300 kg of nitrogen, 100–200 kg of P_2O_5 , and 150–200 kg of K_2O per hectare [Warade and Shinde 1998]. The fertilizer is usually split into three or four doses [Inden and Asahira 1990]. According to Wonneberger et al. [2004] the bunching onion harvested with green tops should have an intensive green colour of leaves and requires $180 \text{ kg N}\cdot\text{ha}^{-1}$, including about $50 \text{ kg N}\cdot\text{ha}^{-1}$ normally present in the soil in mineral forms just before planting. Much lower doses within the range of $60\text{--}80 \text{ kg N}\cdot\text{ha}^{-1}$ are recommended by Skąpski and Dąbrowska [1994]. In China $240 \text{ kg N}\cdot\text{ha}^{-1}$ is considered as the optimum level for Japanese bunching onion cultivated for early spring harvest of blanched pseudostems of this crop [Jiang et al. 2007]. In the greenhouse studies nitrate appeared to be a better source of nitrogen than ammonium which caused a significant suppression of plants growth [Perner et al. 2007]. The knowledge of nitrogen requirement by Japanese bunching onion under Polish climatic conditions is still limited.

The aim of the study was to evaluate the yield and nutritional value of Japanese bunching onion in relation to nitrogen rate and source of this nutrient.

MATERIAL AND METHODS

Field experiment which the aim was to determine the response of Japanese bunching onion to nitrogen fertilization was conducted in 2010–2011 at Piastów Horticultural Experimental Station (long. 17.00 E; lat. 51.05 N) on a sandy clay soil with pH 7.1 and organic matter content 1.8%, Available forms of phosphorus and potassium expressed in 1 dm³ of the soil were raised up to the standard level for onion equal to 80 mg P and 200 mg K by early spring fertilization with triple superphosphate and potassium chloride.

Calcium nitrate [Ca(NO₃)₂ – 15.5% N], ammonium nitrate [NH₄NO₃ – 34% N] and Entec 26 – a mixture of ammonium sulphate and ammonium nitrate with total amount of 26% N, 13% S and addition of DMPP nitrification inhibitor (3,4 dimethylpyrazole phosphate) were used as the sources of nitrogen for Japanese bunching onion. All tested fertilizers were applied in the amounts of 75, 150 and 225 kg N·ha⁻¹ in the single pre-plant doses under harrowing, shortly before planting. The experiment was established in two factorial design in four replications, and plot area amounted 6 m² (1.5 × 4 m).

Seeds of Japanese bunching onion Performer cultivar (Bejo Zaden) were sown into multicell trays filled with a standard peat moss substrate on 25 of January. To each cell containing 54 cm³ of the medium 3–4 seeds were put in, and after emergence the number of seedlings was reduced to two. Transplants produced in the greenhouse during last ten days before planting were hardened in non heated plastic tunnel. Well developed seedlings at the stage of 2–3 leaves were transplanted into field on 7 and 5 of April in the subsequent years, in spacing 30 × 15 cm, which assured the population of 44 plants per 1m². Crop management included hand weeding of plots and sprinkler irrigation of plants in rainfall deficiency periods. In general, weather conditions during all trial periods were favourable for the growth and development of Japanese bunching onion.

Single harvest was conducted on 14 and 15 of June. The whole plants with removed roots and yellowing leaves, and stem diameter > 10 mm were recognized as the marketable yield. Samples of 10 plants from each plot were collected for evaluation such morphological features as mean weight and total length of plant, total length of pseudostem, number of leaves and bulb diameter. In edible parts of plants there were determined the contents of dry matter (by drying at 105°C to the constant weight – PN-90/A-75101/03), total and reducing sugars (Loof-Shoorl method – PN-90A-7501/07), vitamin C (Tillmans method – PN-90/A-7501/11), chlorophyll a + b (spectrophotometric method), and carotenoids (colorimetric method) according to Lichtenthaler and Welburn [1983], volatile oils (distillation method PN – ISO 6571), total N (Kjeldahl method) according to Ostrowska et al. [1991], nitrates expressed by the amount of NO₃-N in f.w. (ion selective electrode, Orions method), P and Mg colorimetrically, Ca and K by photometric method [according to Nowosielski 1974].

The results of the field study and chemical analysis were evaluated statistically using analysis of variance for two factorial design and the least significant differences calculated by Tukey test at $\alpha = 0.05$.

RESULTS AND DISCUSSION

The relations between the tested factors, yield and nutritional value of the crop were similar in both years of the study, and for this reason all the received data are presented as the means for 2010–2011. It was confirmed a high suitability a new pseudostem type Performer cultivar for a single harvest the whole plants supplied for fresh market. Like in the study conducted by Tendaj and Mysiak [2011] it showed little tendency to tillering. The other important features of this cultivars are long pseudostem, and rapid plants growth resulted in average marketable yield of plants after 70 days from transplanting at the range of 15.0–19.5 t·ha⁻¹.

Total yield of Japanese bunching onion was not significantly affected by the source of nitrogen used in the study. Irrespective of N dose, ammonium nitrate and Entec 26 provided similar while calcium nitrate slightly lower total yield of plants (tab. 1). In the case of marketable yield comprising plants with pseudostem diameter > 10 mm this effect was still pronounced but not confirmed statistically. The form on nitrogen fertilizer did not influence the mean weight of plant and all tested morphological features such as height of plant, number of leaves and pseudostem diameter at harvest (tab. 2–3). Also Perner et al. [2007] in pot experiment conducted in the greenhouse conditions did not observe the differences in number of leaves and shoot dry weight in treatments supplied with nitrate and ammonium nitrate forms, while the effect was pronounced in the case of ammonium application. Shoot growth and number of leaves of NH₄⁺ fed plants were significantly suppressed compared to the other mentioned N sources. Similarly Hawkins and George [2001] found that NO₃⁻ fed wheat plants were larger than NH₄⁺ fed ones.

Table 1. Total and marketable yield of Japanese bunching onion as affected by nitrogen fertilization (t·ha⁻¹)

| Kind of fertilizer | Total yield (t·ha ⁻¹) | | | | Marketable yield (t·ha ⁻¹) | | | |
|--|--|------|------|------|--|------|------|------|
| | nitrogen rate (kg N·ha ⁻¹) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 15.5 | 17.4 | 15.9 | 16.3 | 15.2 | 16.6 | 15.0 | 15.6 |
| Ammonium nitrate | 19.0 | 19.2 | 20.0 | 19.4 | 18.6 | 18.7 | 19.5 | 18.9 |
| Entec 26 | 18.3 | 18.9 | 19.7 | 19.0 | 17.9 | 18.3 | 19.2 | 18.5 |
| Mean | 17.6 | 18.5 | 18.5 | 18.2 | 17.2 | 17.9 | 17.9 | 17.7 |
| LSD _{$\alpha=0.05$} for: | A – kind of fertilizer | | | n.s. | | | | n.s. |
| | B – N rate | | | n.s. | | | | n.s. |
| | interaction A × B | | | n.s. | | | | n.s. |

Table 2. Mean weight of plant and pseudostem diameter of Japanese bunching onion as affected by nitrogen fertilization

| Kind of fertilizer | Weight of plant (g) | | | | Pseudostem diameter (mm) | | | |
|----------------------------|--|------|------|------|--------------------------|------|------|------|
| | nitrogen rate (kg N·ha ⁻¹) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 44.7 | 47.2 | 46.8 | 46.2 | 15.7 | 15.7 | 16.0 | 15.8 |
| Ammonium nitrate | 43.0 | 48.4 | 48.5 | 46.6 | 17.3 | 16.8 | 16.6 | 16.9 |
| Entec 26 | 46.5 | 46.5 | 49.6 | 47.5 | 16.9 | 15.9 | 14.8 | 15.9 |
| Mean | 44.7 | 47.4 | 48.3 | 46.8 | 16.6 | 16.1 | 15.8 | 16.2 |
| LSD _{α=0.05} for: | A – kind of fertilizer | | | n.s. | | | | n.s. |
| | B – N rate | | | n.s. | | | | n.s. |
| | interaction A × B | | | n.s. | | | | n.s. |

Table 3. Height of plant and number of leaves in Japanese bunching onion influenced by nitrogen fertilization

| Kind of fertilizer | Height of plant (cm) | | | | Number of leaves | | | |
|----------------------------|--|------|------|------|------------------|-----|-----|------|
| | nitrogen rate (kg N·ha ⁻¹) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 48.2 | 48.7 | 47.5 | 48.1 | 5.1 | 5.0 | 5.0 | 5.0 |
| Ammonium nitrate | 50.5 | 52.0 | 50.6 | 51.0 | 5.1 | 5.3 | 5.2 | 5.2 |
| Entec 26 | 49.9 | 51.5 | 50.4 | 50.6 | 5.1 | 5.2 | 5.1 | 5.1 |
| Mean | 49.5 | 50.7 | 49.5 | 49.9 | 5.1 | 5.2 | 5.1 | 5.1 |
| LSD _{α=0.05} for: | A – kind of fertilizer | | | n.s. | | | | n.s. |
| | B – N rate | | | n.s. | | | | n.s. |
| | interaction A × B | | | n.s. | | | | n.s. |

The increment of nitrogen rate from 75 to 150 and 225 kg·ha⁻¹ did not affect the growth bunching onion, total, and marketable yield as well as height of plants, leaves number and pseudostem diameter. The only exception was a significant enhancement of mean plant weight in treatments fertilized with 150 and 225 kg N·ha⁻¹. In the other studies there was observed higher requirement for this nutrient and significant increment of Japanese bunching yield caused by the application of 166 kg N·ha⁻¹ in comparison to the dose of 100 kg N·ha⁻¹ [Leong 2001]. Similar level of nitrogen fertilization between 130 and 160 kg·ha⁻¹ for onion is recommended by Lorenz and Maynard [1988], while 240 kg·ha⁻¹ by Jiang et al. [2007] in the case of 60 t·ha⁻¹ yield target. Small response to nitrogen in our study could be explained by short only 10 weeks growing period and harvest of plants arranged in early stage of maturation, when majority of them reached 10 mm of stem diameter and provided the yield below 20 t·ha⁻¹. This suggestion can be

confirmed by the data of some other studies with vegetables of short growing period like Swiss chard and spinach, which requirement for N did not exceed $100 \text{ kg} \cdot \text{ha}^{-1}$ [Kołota and Czerniak 2010, Krężel and Kołota 2010].

Both, nitrogen form and rate of this nutrient did not influence the content of dry matter, vitamin C, total sugars and volatile oils (tab. 4–5). It is worth to notice however, that regardless of N feeding level, Japanese bunching onion like in the other trials conducted by Kotlińska and Kojima [2000], Higashio et al. [2007], Tendaj and Mysiak [2007b] appeared to be an abundant source of vitamin C, which content ranged within 28.6 and $32.9 \text{ mg} \cdot 100 \text{ g}^{-1} \text{ f.w.}$

Total chlorophyll content was significantly dependent on nitrogen form and rate (tab. 6). Plants supplied with Entec 26, especially at the doses of 150 and $225 \text{ kg} \cdot \text{ha}^{-1}$ contained higher amounts of this pigment then that with ammonium nitrate, and sub-

Table 4. Dry matter and vitamin C content in Japanese bunching onion influenced by nitrogen fertilization

| Kind of fertilizer | Dry matter (%) | | | | Vitamin C ($\text{mg} \cdot 100 \text{ g}^{-1} \text{ f.w.}$) | | | |
|---------------------------|--|-------|-------|-------|---|-------|-------|-------|
| | Nitrogen rate ($\text{kg N} \cdot \text{ha}^{-1}$) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 10.52 | 10.54 | 10.32 | 10.46 | 31.28 | 28.59 | 32.85 | 30.91 |
| Ammonium nitrate | 10.80 | 10.40 | 10.64 | 10.61 | 30.45 | 29.69 | 31.25 | 30.46 |
| Entec 26 | 10.81 | 10.35 | 9.86 | 10.34 | 29.04 | 29.77 | 29.93 | 29.58 |
| Mean | 10.71 | 10.43 | 10.27 | 10.47 | 30.26 | 29.35 | 31.34 | 30.32 |
| LSD $_{\alpha=0.05}$ for: | A – kind of fertilizer | | | n.s. | n.s. | | | n.s. |
| | B – N rate | | | n.s. | n.s. | | | n.s. |
| | interaction A \times B | | | n.s. | n.s. | | | n.s. |

Table 5. Total sugar and essential oils content in Japanese bunching onion influenced by nitrogen fertilization

| Kind of fertilizer | Total sugar (%) | | | | Essential oils ($\text{ml} \cdot 100 \text{ g}^{-1} \text{ f.w.}$) | | | |
|---------------------------|--|------|------|------|--|------|------|------|
| | nitrogen rate ($\text{kg N} \cdot \text{ha}^{-1}$) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 4.44 | 4.08 | 4.64 | 4.39 | 0.65 | 0.70 | 0.63 | 0.66 |
| Ammonium nitrate | 4.34 | 4.33 | 4.44 | 4.37 | 0.63 | 0.65 | 0.68 | 0.65 |
| Entec 26 | 4.86 | 4.21 | 4.12 | 4.40 | 0.65 | 0.73 | 0.73 | 0.70 |
| Mean | 4.55 | 4.21 | 4.40 | 4.38 | 0.64 | 0.69 | 0.68 | 0.67 |
| LSD $_{\alpha=0.05}$ for: | A – kind of fertilizer | | | n.s. | n.s. | | | n.s. |
| | B – N rate | | | n.s. | n.s. | | | n.s. |
| | interaction A \times B | | | n.s. | n.s. | | | n.s. |

stantially but not proved statistically with those fed by calcium nitrate. More intensive green color of leaves as a result of enhanced chlorophyll content in different species of vegetable crops supplied with fertilizers containing DMPP nitrification inhibitor was observed by Hähndel and Zerulla [2000], Hähndel and Strohm [2001] and Pasda et al. [2001]. Irrespective of the nitrogen form, the increment of its dose from 75 to 150 kg·ha⁻¹ was a beneficial factor for the total chlorophyll amount in edible parts of Japanese bunching onion. Entec 26, beside calcium nitrate appeared to be a favourable source of N for the content of carotenoids.

Table 6. Chlorophyll a + b and carotenoids content in Japanese bunching onion depending on nitrogen fertilization

| Kind of fertilizer Rodzaj nawozu | Chlorophyll a + b (mg·100 g ⁻¹ f.w.) | | | | Carotenoids (µg·100 g ⁻¹ f.w.) | | | |
|-------------------------------------|---|-------|-------|-------|---|--------|--------|--------|
| | nitrogen rate (kg N·ha ⁻¹) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 51.79 | 54.29 | 58.71 | 54.93 | 166.56 | 168.09 | 167.76 | 167.47 |
| Ammonium nitrate | 51.28 | 51.64 | 54.47 | 52.46 | 148.20 | 152.12 | 142.89 | 147.74 |
| Entec 26 | 51.29 | 60.46 | 62.17 | 57.97 | 151.28 | 171.96 | 167.61 | 163.62 |
| Mean | 51.45 | 55.46 | 58.45 | 55.12 | 155.35 | 164.06 | 159.42 | 159.61 |
| LSD _{α=0.05} for: | A – kind of fertilizer | | | 4.04 | | | | 9.77 |
| | B – N rate | | | 3.99 | | | | n.s. |
| | interaction A × B | | | n.s. | | | | n.s. |

Table 7. Nitrogen and nitrates content in Japanese bunching onion depending on nitrogen fertilization

| Kind of fertilizer | Total nitrogen (%) | | | | Nitrates (mg·kg ⁻¹ f.w.) | | | |
|----------------------------|--|------|------|------|-------------------------------------|------|------|------|
| | nitrogen rate (kg N·ha ⁻¹) | | | | | | | |
| | 75 | 150 | 225 | mean | 75 | 150 | 225 | mean |
| Calcium nitrate | 2.03 | 2.43 | 2.37 | 2.28 | 2329 | 3617 | 4075 | 3340 |
| Ammonium nitrate | 1.92 | 2.12 | 2.47 | 2.17 | 2228 | 2651 | 4092 | 2990 |
| Entec 26 | 2.04 | 2.08 | 2.49 | 2.20 | 1932 | 2495 | 2790 | 2406 |
| Mean | 2.00 | 2.21 | 2.44 | 2.22 | 2163 | 2921 | 3652 | 2912 |
| LSD _{α=0.05} for: | A – kind of fertilizer | | | n.s. | | | | 325 |
| | B – N rate | | | 0.22 | | | | 411 |
| | interaction A × B | | | n.s. | | | | n.s. |

Chemical plant analysis showed that tested nitrogen forms did not cause the differences in uptake and total N content but considerably influenced nitrates accumulation in edible parts of Japanese bunching onion (tab. 7). Plants fed with Entec 26 contained

lower amounts of $\text{NO}_3\text{-N}$ in comparison to those supplied with ammonium nitrate. The highest level of this compound comprised plants growing on plots fertilized with calcium nitrate. The decreased $\text{NO}_3\text{-N}$ concentration in Entec 26 and NH_4NO_3 fed plants indicate that intercellular NO_3^- was immediately metabolized in the plants whereas in NO_3^- only fed ones surplus NO_3^- was stored in vacuoles [Perner et al. 2007]. Beneficial effects of Entec 26 use as a source of nitrogen, expressed by the decrease of nitrates accumulation was also observed in many other vegetable species such as celeriac, red beet, lettuce, spinach, cauliflower, leek and carrot [Hähndel and Zerulla 2001, Pasda et al. 2001, Kołota et al. 2007, Kołota and Adamczewska-Sowińska 2007]. This finding is important from the practical point of view taking into account a high tendency of Japanese bunching onion grown for top greens for nitrates accumulation, which in the trial in some treatments exceeded the level $4000 \text{ mg}\cdot\text{kg}^{-1}$ f.w.

Heavy N application at the rates of 150 and $225 \text{ kg}\cdot\text{ha}^{-1}$ caused the increment of nitrogen uptake and nitrates accumulation in edible parts, irrespective of nitrogen form. Like in the trial conducted by Smoleń et al. [2012] no relations were found between the form of nitrogen including calcium nitrate, Entec 26 and ammonium nitrate, and the content of phosphorus, potassium, calcium and magnesium (data not included in the paper). The increased N rate from 75 to $225 \text{ kg}\cdot\text{ha}^{-1}$ had only a beneficial effects on uptake and amount of magnesium in edible parts, which was enhanced from 146 to 184 mg per 1 kg f.w.

CONCLUSIONS

1. Calcium nitrate, ammonium nitrate and Entec 26 appeared to be equally valuable sources for Japanese bunching onion growth and yield.

2. Entec 26, a new concept nitrogen fertilizer containing DMPP nitrification inhibitor caused the increase of total chlorophyll and carotenoids compared to ammonium nitrate, the commonly used nitrogen source. Plants supplied with this fertilizer contained substantially lower amounts of nitrates than with ammonium nitrate and calcium nitrate.

3. The increment of nitrogen rate over $70 \text{ kg}\cdot\text{ha}^{-1}$ was not efficient for yield of Japanese bunching onion and adversely affected the quality of the crop by significant enhancement of nitrates accumulation.

REFERENCES

- Abbey L., Joyce D.C., Aked J., Smith B., 2002. Genotype, sulphur, nutrition and soil type effects on growth and dry matter production of spring onion. *J. Hort. Sci. Biotech.* 77, 340–345.
- Grevsen K., 1989. Effects of sowing dates on different varieties of Welsh onion (*Allium fistulosum* L.) under temperate coastal climate. *Acta Hort.* 242, 319–324.
- Griffith G., Trueman L., Crowter T., Thomas B., Smith B., 2002. Onions – a global benefit to health. *Phytother. Res.* 16, 603–615.

- Hawkins H.J., George E., 2001. Reduced N-15-nitrogen transport through arbuscular mycorrhizal hyphae to *Triticum aestivum* L. supplied with ammonium vs. nitrate nutrition. *Ann. Bot.* 87, 303–311.
- Hähndel R., Strohm M., 2001. Neue stabilisierte N-Dünger: Weitere Erfahrungen. *Gemüse* 6, 13–16.
- Hähndel R., Zerulla W., 2000. Wirkung auf Ertrag und Qualität von Gemüse bei Entec Düngung. *Gemüse* 3, 14–16.
- Hähndel R., Zerulla W., 2001. Effects of ammonium-stabilized N-fertilizer on yield and quality of vegetables. *Acta Hort.* 563, 81–86.
- He Q.W., Su D.S., Zhao D.W., 1989. Shandong famous vegetables. Tsian: Shandong Sci. and Technol. Press, 165–217.
- Helm J., 1956. Die zu Würz – und Speisezwecken Kultivierten Arten der Gattung *Allium* L. Die Kulturpflanze 4, 130–180.
- Higashio H., Hirokane H., Sato E., Tokuda S., Uragami A., 2007. Enhancement of functional compounds in *Allium* vegetables with UV radiation. *Acta Hort.* 744, 357–361.
- Horbowicz M., Kotlińska T., 1998. Zróżnicowanie zawartości flawonoli w niektórych uprawnych i dzikich gatunkach z rodzaju *Allium*. *Zesz. Probl. Post. Nauk. Rol.* 463, 529–537.
- Horbowicz M., Kotlińska T., 2000. Level of flavonols in wild and cultivated *Allium* species. *Acta Hort.* 517, 375–380.
- Inden H., Asahira T., 1990. Japanese bunching onion (*Allium fistulosum* L.). In: Onions and allied crops, vol. 3. Biochemistry, food science, and minor crops, Brewster J.L., Rabinowitch, H.D. (red.). CRC Press, Boca Raton, Florida, 159–178.
- Jiang L.H., Liu Z.H., Chen Q., Lin H.T., Zhang W.J., 2007. Study of the effect of nitrogen on green Chinese spring onion yield and supplying target value. *Plant Nutrit. Fert. Sci.* 13, 890–896.
- Jones H.A., Mann L.K., 1963. Onions and their Allies. Leonard Hill (books) Limited, London.
- Kołota E., Adamczewska-Sowińska K., 2007. Przydatność nawozu Entec 26 w uprawie pora na zbiór wczesny. *Rocz. AR w Poznaniu*, 383, *Ogrodnictwo* 41, 529–532.
- Kołota E., Adamczewska-Sowińska K., Krężel J., 2007. Suitability of Entec 26 as a source of nitrogen for red beet and celeriac. *Veget. Crops Res. Bull.* 67, 47–54.
- Kołota E., Czerniak K., 2010. The effects of nitrogen fertilization on yield and nutritional value of Swiss chard. *Acta Sci. Pol., Hortorum Cultus* 9 (2), 31–37.
- Kotlińska T., Kaniszewski S., Kwiecień A., 2005. Porównanie metod uprawy siedmiolatki (*Allium fistulosum* L.). *Now. Warzyw.* 40, 25–32.
- Kotlińska T., Kojima O., 2000. Siedmiolatka (*Allium fistulosum* L.) jako wartościowa roślina warzywna. *Rocz. AR Pozn.* 323, *Ogrodnictwo* 31 (2), 311–311.
- Krężel J., Kołota E., 2010. The effect of nitrogen fertilization on yield and biological value of spinach grown for autumn harvest. *Acta Sci. Pol. Hortorum Cultus* 9 (3), 183–190.
- Lazić B., Todorovič V., Dardič M., 2002. Effect of production method on earliness and yield of *Allium fistulosum* L. *Acta Hort.* 579, 359–362.
- Leong A.C., 2001. Performance of bunching onion (*Allium fistulosum* L.) cultivars in the lowlands of tropical Malaysia. *Acta Hort.* 555, 99–101.
- Lichtenthaller H.K., Welburn A., 1983. Determination of total carotenoids and chlorophylls a and b of leaf extract in different solvents. *Bioch. Soc. Trans.* 603, 591–592.
- Liu S., He H., Feng G., 2009. Effects of nitrogen and sulphur interaction on growth and pungency of different pseudostem types of Chinese spring onion (*Allium fistulosum* L.). *Sci. Hort.* 121, 12–18.
- Lorenz O.A., Maynard D.N., 1988. Knott's handbook for vegetable growers, third ed. Wiley-Interscience Publication. New York.

- Martinez L.E., Galmarini C.R., Masuelli R.W., 2005. Introgression of *Allium fistulosum* L. into interspecific hybrid backcrosses between *A. fistulosum* L. and *A. cepa* L. *Acta Hort.* 688, 109–115.
- Mysiak B., Tendaj M., 2006. Content of flavonoids in some *Allium* species grown for green bunching. *Veg. Crops. Res. Bull.* 65, 105–110.
- Mysiak B., Tendaj M., 2008. Content of phenolic acids in edible parts of some *Alliums* species grown for the green bunching. *Acta Sci. Pol. Hortorum Cultus* 7 (4), 57–62.
- Nowosielski O., 1974. Metody oznaczania potrzeb nawożenia. PWRiL, Warszawa.
- Ostrowska A., Gawaliński S., Szczubiałkowska Z., 1991. Metody analiz i oceny właściwości gleb i podłoży – katalog. IOŚ, Warszawa.
- Pasda G., Hähndel R., Zerulla W., 2001. Effect of fertilizer with the new nitrification inhibitor DMPP (3,4-dimethylpyrazole phosphate) on yield and quality of agricultural and horticultural crops. *Biol. Fertil. Soil* 34 (2), 85–97.
- Perner H., Schwarz D., Krumbein A., Li X., Eckhard G., 2007. Influence of nitrogen forms and mycorrhizal colonization on growth and composition of Chinese bunching onion. *J. Plant Nutr. Soil Sci.* 170, 762–768.
- Rabinowitch H.D. Currah L., 2002. *Allium* crop science: recent advances. CAB International. Wallingford UK.
- Rubatzky V.E., Yamaguchi M., 1997. *World Vegetables. Principles, production and nutritive values.* Chapman & Hall. New York.
- Skąpski H., Dąbrowska B., 1994. *Uprawa warzyw w polu.* Wyd. SGGW, Warszawa.
- Smoleń S., Sady W., Wierzińska J., 2012. The influence of nitrogen fertilization with Entec 26 and ammonium nitrate on the concentration of thirty-one elements in carrot (*Daucus carota* L.) storage roots. *J. Elementol.* 7 (1), 115–137.
- Steiner D., Milić N., Čanadonović-Brunet J., Kapor A., Štajner M., Poprović B.M., 2006. Exploring *Allium* species as a source of potential medicinal agents. *Phytother. Res.* 20, 58584.
- Tendaj M., Mysiak B., 2007a, Plonowanie cebuli siedmiolatki (*Allium fistulosum* L.) w zależności od terminu sadzenia rozsady i stosowania płaskich okryć. *Annales UMCS Sec. Horticultura* 17 (2), 5–10.
- Tendaj M., Mysiak B., 2007b, Usefulness of Japanese bunching onion (*Allium fistulosum* L.) for forcing in greenhouse. *Acta Agrobot.* 60 (1), 143–146.
- Tendaj M., Mysiak B., 2010. Charakterystyka wzrostu cebuli siedmiolatki (*Allium fistulosum* L.) uprawianej z siewu i rozsady. *Konf. Nauk. pt. „Proekologiczna uprawa warzyw – problemy i perspektywy”.* Siedlce 24–25 czerwca, 163–165.
- Tendaj M., Mysiak B., 2011. Growth characteristic of Welsh onion (*Allium fistulosum* L.) grown from seeds and transplants. *Folia Hort.* 23, 1, 3–8.
- Wang D., Gao J., Liu G., 2005. General situation of *Allium* crops in China. *Acta Hort.* 688, 327–332.
- Warade S.D., Shinde K.G., 1998. *Handbook of vegetable science and technology: production, composition storage and processing.* Ed. by Salunkhe D.K. and Kadam S.S., Marcel Dekker. Inc. New York.

REAKCJA CEBULI SIEDMIOLATKI (*Allium fistulosum* L.) NA NAWOŻENIE AZOTEM

Streszczenie. Wprowadzenie do uprawy nowych odmian siedmiolatki o wydłużonej łodydze rzekomej może stanowić alternatywę dla uprawianej powszechnie cebuli zwyczajnej na wczesny zbiór pęczkowy. Jak dotąd brak jest dokładnych informacji na temat potrzeb nawozowych tego warzywa w stosunku do azotu. W dwuczynnikowym doświadczeniu polowym porównywano skuteczność nawożenia saletrą wapniową, saletrą amonową i nawozem Entec 26 zawierającym DMPP jako inhibitor nityfikacji w dawkach przedwegetacyjnych w wysokości 75, 150 i 225 kg·ha⁻¹. Rozsadę wyprodukowaną w szklarni sadzono na miejsce stałe na początku kwietnia, zaś zbiór roślin wykonywano po 10 tygodniach uprawy. Ocenie poddano plon oraz wartość biologiczną roślin wyrażoną zawartością suchej masy, witaminy C, chlorofilu ogółem, karotenoidów, olejków lotnych, N ogólnego N-NO₃(V), P K Ca i Mg. Badania wykazały, że testowane nawozy miały równorzędną przydatność jako źródło azotu dla cebuli siedmiolatki. Użycie nawozu Entec 26 zapewniło większą zawartość chlorofilu i karotenoidów w roślinach w stosunku do powszechnie stosowanej w nawożeniu saletry amonowej. Innym korzystnym efektem stosowania tego nawozu był istotny spadek zawartości azotanów w częściach jadalnych. Wzrost przedwegetacyjnie stosowanej dawki azotu z 75 do 150 i 225 kg·ha⁻¹ nie miał wpływu na wielkość plonu, przyczynił się jednak do znacznego wzrostu poziomu akumulacji azotanów w roślinach.

Słowa kluczowe: forma azotu, dawka N, plon, wartość odżywcza roślin

ACKNOWLEDGEMENT

Presented paper contains the research data obtained within the framework of grant no: 4454/B/PO/2010/38 financed by the Polish Ministry of Science and Higher Education.

Accepted for print: 19.07.2012