

PARASITIC WASPS OF THE *PIMPLINAE* SUBFAMILY (*Hymenoptera*, *Ichneumonidae*) OF AGRICULTURAL LANDSCAPE REFUGIUM HABITATS IN CENTRAL WIELKOPOLSKA

Hanna Piekarska-Boniecka, Barbara Wilkaniec,
Ewa Dolańska-Niedbała

University of Life Sciences in Poznań

Abstract. The research was carried out in the years 1991–1994 in agricultural landscape of medium mosaic degree in the area of Łęczycza. It covered parasitoids of the *Pimplinae* subfamily (*Hymenoptera*, *Ichneumonidae*), inhabiting refugium habitats such as shrubberies, field borders, roadsides and forest edges. The method used was scooping in order to catch imagos. 48 species were found, which made up 36.1% of this subfamily's domestic fauna and 63.1% of those reported for Wielkopolska. Dominant species were *Endromopoda detrita* (Holmgr.) ($D = 19.5\%$), *Itoplectis maculator* (F.) ($D = 11.5\%$), *I. alternans* (Grav.) ($D = 11.3\%$), *Pimpla contemplator* (Muell.) ($D = 9.2\%$) and *Zaglyptus multicolor* (Grav.) ($D = 7.1\%$). The greatest species diversity was reported from the forest edge (34 species), while the highest number was found in the shrubberies (268 individuals). The greatest stability of species composition was found in the shrubbery communities. The greatest species similarity between communities was reported for the shrubberies and the forest edge ($SM = 53.5\%$). The highest similarity in terms of quantity occurred between the groups on the field border and road edge ($Re = 64.5\%$).

Key words: parasitoid, *Pimplinae*, refugium habitats, agricultural landscape, Wielkopolska

INTRODUCTION

Refugium habitats in an agricultural landscape are a basic element increasing agrocenoses' species diversity. In this way they positively influence their stability, frequently deciding on the distribution and development of various beneficial species, also parasitoids. The impact of refugium habitats on parasitoid entomofauna has been the subject of foreign studies [eg. Landris and Haas 1992, Altieri et al. 1993, Marino and

Corresponding author – Adres do korespondencji: Hanna Piekarska-Boniecka, Department of Entomology, University of Life Sciences in Poznań, ul. Dąbrowskiego 159, 60-594 Poznań, Poland, e-mail: boniecka@au.poznan.pl

Landris 1996, 2000]. Polish resources do not provide many resources on the subject. The influence of landscape structure on the occurrence of parasitoids of the *Ichneumonidae* family, which are a decisive factor in biocenose-controlling processes in agroecosystems, was presented in studies by Strawiński [1957], Kościelska [1959], Piekarska-Boniecka [1999, 2005] and Piekarska-Boniecka and Wilkaniec [2001, 2006]. Thus it was decided to start research on parasitic entomofauna infesting agroecosystems of various mosaics degree.

The research was aimed at defining the quality-quantity composition of the *Pimplinae* subfamily parasitoids occurring in refugium habitats, such as woodlots, field borders, roadsides and forest edges, in agricultural landscape of medium mosaic degree in Wielkopolska.

MATERIAL AND METHODS

The research was conducted in the years 1991–1994 in central Wielkopolska, on cultivated fields located between the towns of Puszczykowo and Łęczyca (UTM: XT29), which constituted agricultural landscape of complex type, of medium mosaic degree. The share of non-cultivated habitats was 7.6%. The fields up to 30 ha surface area were situated on the flat top of the so-called Wirska Moraine and bordered with a poor form of Central-European oak-hornbeam complex (*Galio silvatici-Carpinetum*), located in Wielkopolska National Park.

Parasitic wasps were caught from May to October of each year of study using the scooping method. The catch was conducted on the following sites: shrubberies, field borders, roadside and forest edge. 10 samples per month were collected per site. Insects caught in 25 scoop strokes were assumed as one sample.

Phytosociological characteristics of the sites was created on the basis of the study by Matuszkiewicz [2001], while the terminology of vascular plants was adopted according to the list of taxa provided in the study by Mirek et al. [2002].

The plant cover of habitats – places of the research was the following:

– shrubberies – 150 m long, made up by *Pruno-Crataegetum*, with nitrophilous edge and ruderal communities on both sides, including plots of *Convolvulo-Agrophyretum*,

– field border – 250 m long, overgrown with graminoids (*Convolvulo-Brometum inermis*, *Convolvulo-Agrophyretum* and *Arrhenatheretum medioeuropaeum*) with some species of exothermic grass cover of *Festuco-Brometea* class and thermophilous edges with *Trifolio-Geranietea*,

– midfield roadside – 300 m long, the most numerous plant community was a group of *Convolvulo-Agrophyretum*. Under the cover of trees and shrubs edge groupings occurred. Among trees the dominants were the sycamore maple (*Acer pseudoplatanus* L.), norway maple (*Acer platanoides* L.), common ash (*Fraxinus excelsior* L.) and sessile oak (*Quercus petraea* (Matuschka) Lie),

– forest edge – 300 m long, it was the zone between poor form of central-European oak-hornbeam complex and fields. Along the forest edge ran a dust road overgrown with patches of trampled *Lolio-Plantaginetum* community. In forest complex the pine

(*Pinus sylvestris* L.) was dominant. Edge thicket groupings included midfield thickets (*Pruno- Crataegum*), with quite numerous elder plants (*Sambucus nigra* L.). Also herbaceous vegetation and ruderal groupings were present.

On fields, in direct neighbourhood of shrubberies, winter rye, spring barley, winter rape and winter wheat were cultivated. The field border divided fields on which winter and spring wheat, oats, spring and winter rye were cultivated. Arable lands adjacent to the road edges were earmarked for winter wheat, winter rye, maize, oat and winter rape. The forest edge bordered on fields where winter rape, winter and spring wheat and spring barley were cultivated.

The groupings of parasitoids caught in particular habitats were characterised on the basis of the following biocenotic indices: Shannon's diversity index (H') [Shannon and Weaver 1963], Pielou's evenness index [1966] (J') and Margalef's richness index [1958] (d). The stability of species structure of parasitoids communities was analysed on the basis of modified index of community homogeneity in time – Schwerdtfeger's index [1975] (Kj). Parasitoid communities were compared in terms of quality with Marczewski-Steinhaus' index (MS) [Marczewski and Steinhaus 1959] and in terms of quantity with Renkonen's number [1938] (Re).

RESULTS

In refugium habitats of agricultural landscape of Łęczycza area 960 samples were collected in the years 1991–1994. 806 *Pimplinae* specimens of 48 species were caught (Tab. 1, 2). The species reported made up 36.1% of domestic fauna of this subfamily and 63.1% of those reported for Wielkopolska. Eudominants include *Endromopoda detrita* ($D = 19.5\%$), *Itopectis maculator* ($D = 11.5\%$) and *I. alternans* ($D = 11.3\%$), while dominants were: *Pimpla contemplator* ($D = 9.2\%$) and *Zaglyptus multicolor* ($D = 7.1\%$).

Species diversity in particular habitats was very similar. On the forest edge, in shrubberies and on the roadside 34 to 31 species were found (Tab. 1, 2). Only on the field border slightly fewer of them were caught: 25. Such shape of diversity was confirmed by the values of Margalef's richness index (d) and Shannon's diversity index (H'), with the highest value at 14.72 and 2.37 on the forest border, and the lowest at 10.42 and 2.1 on the field border (Tab. 2).

The dominant structure of the *Pimplinae* communities in particular habitats was also very similar. Besides species dominating in all the habitats, taking them as a whole, in shrubberies *Scambus inanis* ($D = 5.7\%$) also occurred numerously. Similarly, on the field border *Scambus brevicornis* ($D = 8.9\%$) and *Pimpla spuria* ($D = 8.4\%$) were also abundant, as was the case on the roadside with *P. rufipes* ($D = 7.4\%$) and *P. spuria* ($D = 7.4\%$), while on the forest edge the numerous species was *Polysphincta tuberosa* ($D = 6.9\%$).

Dominant *Pimplinae* species may control the number of phytophages occurring in agricultural cultivations belonging to genera *Lepidoptera*, *Coleoptera*, *Hymenoptera* and *Diptera*. Among particularly important species due to their biocenotic functions are *Endromopoda detrita*, parasitoids of *Cephus pygmaeus* (L.) and species of the *Itopectis*

genus, which are listed among polyphages. Among dominants was one species of *Polysphincta tuberosa*, which is a parasitoid of predators of *Arachnida*.

The *Pimplinae* communities were characterised with similar mean values of Schwerdtfeger's index (Kj) at the same time, describing yearly changes in species composition in four-year long research. The Kj index values fluctuated between 42.9% and 51.6% (Tab. 3). This means that the communities quality composition reported each year in particular habitats was generally identical in about 50% in the research periods. It was the most stable in shrubberies and the most fluctuating on forest edge.

Species composition of *Pimplinae* communities reported in particular refugium habitats did not show significant similarity, as Marczewski-Steinhaus index (SM) achieved values from 43.9% to 53.5% (Tab. 4). The highest species similarity was between the communities in shrubberies and on the forest edge ($SM = 53.5\%$), while the lowest between *Pimplinae* found on field border and forest edge ($SM = 43.9\%$).

The number of *Pimplinae* specimens found in particular habitats was clearly diversified (Tab. 1, 2). Definitely the most specimens were caught in shrubberies, where 268 specimens were found. The lowest number was found on the road edge, where 163 specimens were caught. The road and forest edges were habitats with the lowest number of *Pimplinae* specimens caught.

The distribution of *Pimplinae* occurrence in particular habitats was even. The values of Pielou's index (J') were similar and at the same time mean and obtained the values between 0.46 and 0.51 (Tab. 2). Such a level of J' values proves that the quantity structure of communities was fully developed. The communities had species with gradually diminishing number and a large number of species with very low number.

The quantity composition of *Pimplinae* communities in particular habitats was varied. Renkonnen's number (Re) showed values from 50.3% to 64.5% (Tab. 4). The communities most similar in terms of number were found on the field border and roadside ($Re = 64.5\%$), while the biggest differences were found in shrubberies and on the forest edge ($Re = 50.3\%$).

DISCUSSION

In the years 1991–1994 in agricultural landscape of medium mosaic degree of in central Wielkopolska 48 *Pimplinae* species were found, which constitute 63.1% of species reported for the area. This proves that refugium habitats in agricultural landscape are places of occurrence of parasitic wasps of *Pimplinae* subfamily, and that is why they can control the number of phytophagous species connected with agricultural cultivations. Those biocenotic relations corroborated research results in Wielkopolska as presented in the studies by Piekarska-Boniecka [2005] and Piekarska-Boniecka and Wilkaniec [2006]. Positive influence of varied structure of agricultural landscape of western Wielkopolska on the occurrence of entomofages, including parasitoids of *Ichneumonidae* subfamily, was also confirmed in the study by Karg [1989]. Positive impact of refugium habitats in agricultural landscape on species diversity and number of parasitic wasps was also presented in studies by Bilewicz-Pawińska and Pankanin-Franczyk [1995] as well as Barczak et al. [2000, 2002].

Research confirmed the dominant structure of *Pimplinae* caught in all the refugium habitats taken as a whole, as those species had previously been reported as dominant in those habitats by Piekarska-Boniecką [2005] and Piekarska-Boniecka and Wilkaniec [2006].

The research showed that the species stability of *Pimplinae* communities was lower on particular sites than reported in the study by Piekarska-Boniecka [2005]. Schwerdtfeger's index (K_j) showed lower values. This proves that species composition of the groupings undergoes clear changes in subsequent years of study.

The research presented the largest species diversity of *Pimplinae* on the forest edge, and the highest number in shrubberies. This was not confirmed by research by Piekarska-Boniecka [2005], who proved that on the field border and roadside the *Pimplinae* community was the most diversified in terms of species, while on the forest edge parasitic wasps were the most numerous. This proves that parasitoids move in consequent years within those habitats.

The research proved the largest species similarity of *Pimplinae* communities found in shrubberies and on the road edge. It corroborated the value of Marczewskie-Steinhaus index (SM) for those groupings, as previous studies by Piekarska-Boniecka [2005] defined it at the same level.

CONCLUSIONS

Refugium habitats in agricultural landscape of medium mosaics degree of in central Wielkopolska, with 7.6% share of non-arable habitats, constitute attractive habitats for parasitoids of the *Pimplinae* subfamily (*Hymenoptera, Ichneumonidae*). The forest edge is the habitat attracting the most species of parasitic wasps, while in midfield thickets the number is the greatest. Agroecosystems should be organised in such a way that the share of refugium habitats for beneficial entomofauna in the form of shrubberies, field borders, roadside and forest edges is about 10% of all the agricultural area. Such a structure of agricultural landscape ensures parasitoids appropriate conditions for living, shelter, breeding and enables their dispersion onto neighbouring cultivations.

REFERENCES

- Altieri M. A., Cure M.A., Garcia M.A., 1993. The role and enhancement of parasitic *Hymenoptera* biodiversity in agroecosystems. [In:] LaSalle J., Gauld I.D. (eds.), *Hymenoptera and biodiversity*. C.A.B. International, Oxon, 257–276.
- Barczak T., Kaczorowski G., Bennewicz J., Krasicka-Korczyńska E., 2000. Znaczenie zarośli śródpolnych jako rezerwarów naturalnych wrogów mszyc. Wyd. AT-R Bydgoszcz: 147 pp.
- Barczak T., Bennewicz J., Kaczorowski G., 2002. Zarośla śródpolne jako rezerwar bioróżnorodności afidofagów. In: Banaszak J. (eds.), *Wyspy środowiskowe. Bioróżnorodność i próby typologii*. Wyd. Akad. Bydg., Bydgoszcz, 127–155.
- Bilewicz-Pawińska T., Pankanin-Franczyk M., 1995. Wpływ czynników środowiska na parazytoidy (*Hymenoptera*) w agroekosystemach. *Wiad. Entomol.*, 14 (2), 103–111.

- Karg J., 1989. Zróżnicowanie liczebności i biomasy owadów latających krajobrazu rolniczego zachodniej Wielkopolski. Roczn. AR Pozn. Rozpr. Nauk. 188, 78 pp.
- Kościelska M. K., 1959. Śródpolne gasieniczniki Polski Zachodniej (*Hym.*, *Ichneumonidae*). Pol. Pismo Entomol., 29, 179–194.
- Landris D.A., Haas M., 1992. Influence of landscape structure on abundance and within-field distribution of *Ostrinia nubilalis* Huebner (*Lepidoptera*, *Pyralidae*) larval parasitoids in Michigan. Environ. Entomol. 21, 409–416.
- Marczewski E., Steinhaus H., 1959. O odległości systematycznej biotopów. Zast. Mat., 4, 195–203.
- Margalef R., 1958. Information theory in ecology. General Systems, 3, 36–71.
- Marino P.C., Landis D.A., 1996. Effect of landscape structure on parasitoid diversity and parasitism in agroecosystems. Ecol. Appl. 6, 1, 276–284.
- Marino P.C., Landis D. A., 2000. Parasitoid community structure implications for biological control in agricultural landscape. [In:] Ekbom B., Irwin M.E., Robert Y. (eds.), Interchanges of insect between agricultural and surrounding landscape. Kluwer Acad. Publ., Dordrecht: 183–193.
- Matuszkiewicz W., 2001. Przewodnik do oznaczania zbiorowisk roślinnych Polski. PWN Warszawa, 536 pp.
- Mirek Z., Piękoś-Mirkowa H., Zając A., Zając M., 2002. Krytyczna lista roślin naczyniowych Polski. Wyd. Kraków, Inst. Bot. W. Szafera PAN, Ser. Różnorodność Biologiczna Polski I, 442 pp.
- Piekarska-Boniecka H., 1999. *Pimplinae* i *Diacritinae* (*Hymenoptera*, *Ichneumonidae*) wybranych środowisk krajobrazu rolniczego okolic Poznania. Cz. 1. Wiad. Entomol., 18 (2), 77–84.
- Piekarska-Boniecka H., 2005. Dynamika zgrupowań *Pimplinae* (*Hymenoptera*, *Ichneumonidae*) w krajobrazie rolniczym środkowej Wielkopolski. Roczn. AR Pozn., Rozpr. Nauk. 366: 204 pp.
- Piekarska-Boniecka H., Wilkaniec B., 2001. *Pimplinae* and *Diacritinae* (*Hymenoptera*, *Ichneumonidae*) occurring in refuge habitats of an agricultural landscape of Poznań environs. Roczn. AR Pozn., 339, Ogródnictwo 34, 59–65.
- Piekarska-Boniecka H., Wilkaniec B., 2006. *Pimplinae*, *Diacritinae*, *Poemeniinae* and *Rhyssinae* parasitoid subfamilies (*Hymenoptera*, *Ichneumonidae*) of refugium habitats in agricultural landscape of central Wielkopolska. Roczn. AR Pozn. 379, Ogródnictwo 40, 33–41.
- Pielou E. C., 1966. Shannon's formula as a measure of specific diversity; its use and misuse. Am. Nat., 100, 463–465.
- Renkonen O., 1938. Statistisch-ökologische Untersuchungen "über die terrestrische Käferwelt der finnischen Bruchmoore". Zool. Soc. Zoo. Bot. Fenn., 6, 1–226.
- Schwerdfeger F., 1975. Ökologie der Tiere. Synökologie. Parey Verlag, Hamburg, Berlin, 451 pp.
- Shannon C. E., Weaver W., 1963. The mathematical theory of communication. Urbana.
- Strawiński K., 1957. Zmiany zachodzące w entomofaunie przy zakładaniu zadrzewień śródpolnych. Pol. Pismo Entomol., Ser. B 4 (4), 49–72.

PASOŻYTNICZE BŁONKÓWKI Z PODRODZINY *Pimplinae* (Hymenoptera, Ichneumonidae) ŚRODOWISK OSTOJOWYCH KRAJOBRAZU ROLNICZEGO ŚRODKOWEJ WIELKOPOLSKI

Streszczenie. Badania prowadzono w latach 1991–1994 w krajobrazie rolniczym o średnim stopniu mozaikowości w okolicach Łęczycy. Badaniami objęto parazytoidy z podrodziny *Pimplinae* (Hymenoptera, Ichneumonidae), zasiedlające środowiska ostojoye w postaci zakrzewień, miedz, przydroży i skrajów lasu. Wykorzystano metodę czerpakowania w celu odłowu imagines. Stwierdzono występowanie 48 gatunków, które stanowiły 36,1% fauny krajowej tej podrodziny i 63,1% wykazanych z Wielkopolski. Gatunkami dominującymi były *Endromopoda detrita* (Holmgr.) ($D = 19,5\%$), *Itoplectis maculator* (F.) ($D = 11,5\%$), *I. alternans* (Grav.) ($D = 11,3\%$), *Pimpla contemplator* (Muell.) ($D = 9,2\%$) i *Zaglyptus multicolor* (Grav.) ($D = 7,1\%$). Wykazano najwyższe zróżnicowanie gatunkowe na skraju lasu (34 gatunki), i najwyższą liczebność w zakrzewieniach (268 osobników). Stwierdzono największą stabilność składu gatunkowego zgrupowań występujących w zakrzewieniach. Wykazano największe podobieństwo gatunkowe pomiędzy zgrupowaniami stwierdzonymi w zakrzewieniach i na skraju lasu ($SM = 53,5\%$). Stwierdzono największe podobieństwo w kategoriach ilościowych pomiędzy zgrupowaniami występującymi na miedzy i przydrożu ($Re = 64,5\%$).

Key words: parazytoid, *Pimplinae*, środowiska ostojoye, krajobraz rolniczy, Wielkopolska

Accepted for print – Zaakceptowano do druku: 4.11.2008